

Evaluating Patients For Secondary Syphilis (P1/3)

*SEXUAL HISTORY, RISK ASSESSMENT & PHYSICAL EXAM

Sexual History, Risk Assessment (past year):

- gender of partners
- number of partners (new, anonymous, serodiscordant HIV status, exchange of sex for drugs or money)
- types of sexual exposure
- recent STDs; HIV serostatus
- substance abuse
- condom use

Physical Exam

- oral cavity
- lymph nodes
- skin
- palms & soles
- neurologic
- genitalia/pelvic
- perianal

History of syphilis

prior syphilis (last serologic test & last treatment)

†DIAGNOSTIC ISSUES IN SECONDARY SYPHILIS

RPR/VDRL

- ~100% sensitive in secondary syphilis
- Tests must be quantified to the highest titer & titer on the day of treatment must be used to assess treatment response
- Always use the same testing method (RPR or VDRL) in sequential testing; cannot compare titer from the two tests
- Tests lack specificity (biologic false positive); all reactive tests need to be confirmed by a treponemal test for syphilis diagnosis
- Prozone Reaction: false negative RPR or VDRL from excess antibody blocking the antigen-antibody reaction
 - ~1% of secondary syphilis cases
 - Request lab to dilute the serum to at least 1/16 to rule out

TREATMENT & FOLLOW-UP

‡Treatment of Secondary Syphilis

Recommended Regimen

- Benzathine Penicillin G 2.4 million units IM x 1

Alternative Regimens for Penicillin Allergic Non-Pregnant Patients:

efficacy not well established & not studied in HIV+; close follow-up essential:

- Doxycycline 100 mg po bid x 2 weeks or
- Tetracycline 500 mg po qid x 2 weeks or
- Ceftriaxone 1gm IM or IV qd x 10-14 d

See CDC 2010 STD Treatment Guidelines:

www.cdc.gov/std/treatment/2010/default.htm

& California STD Treatment Guidelines Grid:

www.stdhivtraining.org/resource.php?id=15&ret=clinical_resources


** Follow-Up To Assess Treatment Response

- 1-2 weeks & 1 month: clinical follow-up
- 3, 6, 9, 12, 24 months: serologic follow-up for HIVinfected
- 6, 12 months: serologic follow-up for HIV negative
- Treatment failure: failure of titer to decline fourfold within 6-12 months from titer at time of treatment

§REPORTING & PARTNER MANAGEMENT

- All syphilis cases or suspected cases must be reported to the local health department within one working day of diagnosis
- Local health departments will assist in partner notification & management
- Contact Number at Local Health Department

Evaluating Patients For Secondary Syphilis (P2/3)




***, †, ‡, §, ** see color coded boxes**

1. All patients with suspected syphilis should be tested for HIV infection & screened for other STDs. Repeat HIV testing of patients with secondary syphilis 3 months after the first HIV test, if the first test is negative

To Order Additional Copies

see the online version of the Secondary Syphilis Algorithm on the clinical resources page of the CA STD/HIV PTC website: <http://www.stdhivtraining.org>



CALIFORNIA
STD/HIV PREVENTION
TRAINING CENTER

Acknowledgements

The California STD/HIV Prevention Training Center thanks the Medical Directors from the National Network of Prevention Training Centers, The California STD Controllers Association and the Division of STD Prevention of the Centers for Disease Control and Prevention for their assistance in preparing this document. Revised 4/2011

