Essential Sexual Health Questions to Ask Adults

Ask all of your adult patients the questions on this card to start the conversation and to begin taking a thorough sexual history. For more questions to assess risk, see Table 1 of "Sexual Health and Your Patients: A Provider's Guide."

Ask at Least Annually Have you been Have you ever been NO sexually active in the sexually active? last year? YES YES → What types of sex → What types do you have (oral, of sex did you vaginal, anal)? have (oral, NO vaginal, anal)? Continue → With men, women, or both? → With men, with medical women, or both? history. In the past 12 months, how many How many sexual sexual partners have partners have you had? you had?

Ask Older Adults

Has sex changed for you? If so, how?

Ask at least once, and update as needed. Gender identity and sexual orientation can be fluid.

- 1. What do you consider yourself to be?
 - A. Lesbian, gay, or homosexual
 - B. Straight or heterosexual
 - c. Bisexual
 - D. Another (please specify)
 - F. Don't know
- 2. What is your current gender identity?
 - A. Male
 - в. Female
 - c. Transgender man
 - D. Transgender woman
 - E. Neither exclusively male nor female (e.g. non-binary or nonconforming)
 - F. Another (please specify)
 - G. Decline to answer
- 3. What sex were you assigned at birth?
 - A. Male
 - **B** Female
 - c. Decline to answer



Recommended Preventive Sexual Health Services for Adults

Service	Females			Males			Transgender
	18-64	65+	Pregnant	18-64	65+	MSM	Individuals
STI Counseling	√ a	√ a	✓a	✓a	✓a	√a	√a
Contraceptive Counseling	✓		✓	✓	✓		✓
Cervical Cancer Screening	✓b	√b	✓b				✓c
Chlamydia Screening	✓d	✓d	✓d	✓e		√f	√ a
Gonorrhea Screening	✓d	✓d	✓d			√g	✓a
HIV Testing	✓	√a	✓	✓	√a	✓	✓
Syphilis Screening	√h	√h	✓	√h	√h	✓	√h
Hepatitis B Screening	√i	√i	✓	√i	√i	✓	√i
Hepatitis C Screening	√ jk	√ jk	√ j	√ jk	√ jk	√ jk	√ jk
Hepatitis A Vaccine	✓I	√I	√ 1	✓I	✓I	✓	√ I
Hepatitis B Vaccine	√m	√m	√m	✓m	√m	✓	√ m
HPV Vaccine	√n			✓n		√n	√n
PrEP	✓*	✓*	✓*	✓*	✓*	✓*	✓*

* = HIV-negative and at substantial risk for HIV infection (sexual partner with HIV, injection drug user, recent bacterial STI, high number of sex partners, commercial sex worker, lives in high-prevalence area or network)

- **a** = At increased risk: inconsistent condom use, multiple partners, partner with concurrent partners, current STI, or history of STI within a year
- **b** = Aged 21 to 65 or when adequate screening history has been established
- **c** = FTM transgender patients who still have a cervix according to guidelines for non-transgender women
- **d** = Sexually-active women aged <25; women aged ≥25 at increased risk
- e = Young adult males in high-prevalence communities or settings
- **f** = Screen for urethral infection if insertive anal sex in preceding year; rectal infection if receptive anal sex in preceding year
- g= Screen for urethral infection if insertive anal sex in preceding year; rectal infection if receptive anal sex in preceding year; pharyngeal infection if receptive oral sex in preceding year
- h = HIV-positive; at increased risk: exchange sex for drugs or money; in high prevalence communities
- **i** = At risk: HIV-positive, unprotected sex, share needles, family member or sexual partner infected with HBV; born in a HBV-endemic country; born to parents from a HBV-endemic country
- j = HIV-positive, history of injection or intranasal drug use or incarnation; blood transfusion prior to 1992
- k = Born between 1945 and 1965 (at least once)
- I = Use illicit drugs; have chronic liver disease; receive clotting factors; travel to HAV-endemic countries; wish to be vaccinated
- **m** = At risk: multiple partners, share needles, family member or sexual partner infected with HBV
- n = Women and men aged ≤45

For more information, visit: national coalition for sexual health.org