

PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Male Adolescent Sexual and Reproductive Health Care

Arik V. Marcell, Charles Wibbelsman, Warren M. Seigel and the Committee on
Adolescence

Pediatrics; originally published online November 28, 2011;
DOI: 10.1542/peds.2011-2384

The online version of this article, along with updated information and services, is
located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/early/2011/11/22/peds.2011-2384>

PEDIATRICS is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. PEDIATRICS is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2011 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 0031-4005. Online ISSN: 1098-4275.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™





CLINICAL REPORT

Male Adolescent Sexual and Reproductive Health Care

abstract

FREE

Male adolescents' sexual and reproductive health needs often go unmet in the primary care setting. This report discusses specific issues related to male adolescents' sexual and reproductive health care in the context of primary care, including pubertal and sexual development, sexual behavior, consequences of sexual behavior, and methods of preventing sexually transmitted infections (including HIV) and pregnancy. Pediatricians are encouraged to address male adolescent sexual and reproductive health on a regular basis, including taking a sexual history, performing an appropriate examination, providing patient-centered and age-appropriate anticipatory guidance, and delivering appropriate vaccinations. Pediatricians should provide these services to male adolescent patients in a confidential and culturally appropriate manner, promote healthy sexual relationships and responsibility, and involve parents in age-appropriate discussions about sexual health with their sons. *Pediatrics* 2011;128:e1658–e1676

INTRODUCTION

During adolescence, a number of changes occur for boys, including the physical, psychological, and social changes associated with puberty, and the majority of male adolescents report the initiation of sexual behavior.¹ Many of these events, including sexual initiation, are associated with preventable consequences that can lead to significant morbidity and mortality.² During this same time period, the number of health visits typically declines, particularly among older male adolescents, and there is a shift from routine to more time-limited acute visits.³

For health care providers, including primary care providers and pediatricians, who care for male adolescents, issues of puberty and sexuality are areas that should be commonly addressed with the male patient and his family.⁴ Even after the release of the American Medical Association's *Guidelines for Adolescent Preventive Services* (GAPS)⁵ and *Bright Futures*,⁶ which recommend preventive health services for adolescents, few improvements have been observed in the counseling of male teenagers regarding the prevention of sexually transmitted infections (STIs) or HIV infection.^{7,8} Furthermore, data from outpatient ambulatory medical records show that primary care providers are 3 times more likely to take sexual health histories from female than male patients and twice as likely to counsel female patients on the use of condoms.^{9,10} Thus, it is important for primary care providers to have an understanding of what sexual/reproductive health care means for the male adolescent.

Addressing male teenagers' sexual/reproductive health includes but is not limited to preventing STIs and HIV. The 1994 Cairo United Nations

Arik V. Marcell, MD, MPH, Charles Wibbelsman, MD, Warren M. Seigel, MD, and the Committee on Adolescence

KEY WORDS

male sexual health, male reproductive health, male adolescents, male puberty

ABBREVIATIONS

STI—sexually transmitted infection
SMR—sexual maturity rating
AAP—American Academy of Pediatrics
NSFG—National Survey of Family Growth
CI—confidence interval
HPV—human papillomavirus
CDC—Centers for Disease Control and Prevention
USPSTF—US Preventive Services Task Force

This document is copyrighted and is property of the American Academy of Pediatrics and its Board of Directors. All authors have filed conflict of interest statements with the American Academy of Pediatrics. Any conflicts have been resolved through a process approved by the Board of Directors. The American Academy of Pediatrics has neither solicited nor accepted any commercial involvement in the development of the content of this publication.

The guidance in this report does not indicate an exclusive course of treatment or serve as a standard of medical care. Variations, taking into account individual circumstances, may be appropriate.

www.pediatrics.org/cgi/doi/10.1542/peds.2011-2384

doi:10.1542/peds.2011-2384

All clinical reports from the American Academy of Pediatrics automatically expire 5 years after publication unless reaffirmed, revised, or retired at or before that time.

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2011 by the American Academy of Pediatrics

International Conference on Population and Development and the World Health Organization defined sexual/reproductive health as “a state of physical, mental and social well-being and not merely the absence of disease, dysfunction or infirmity, in all matters relating to the reproductive system, its functions and its processes.”^{11,12} Sexual health also requires a positive and respectful approach to sexuality and sexual relationships. People should be able to have pleasurable and safe sexual experiences free of coercion, discrimination, or violence. Men, along with women, have the right to be informed and have access to safe, effective, affordable, and acceptable methods of family planning of their choice and the right of access to appropriate health care services.

For health care providers, goals for male adolescents’ sexual/reproductive health beyond the prevention of STIs, HIV infection, unwanted pregnancy, and reproductive health-related cancers should include promoting sexual health and adolescent development, healthy intimate relationships and responsible behavior, and responsible fatherhood as well as reducing problems related to sexual dysfunction and infertility.¹³ Health care providers require the knowledge, skills, and confidence to screen and examine male adolescents and discuss with them and provide appropriate education about the scope of sexual/reproductive health care for male adolescents.

The goals of this report are to:

1. describe key components of male adolescents’ sexual/reproductive health;
2. explain how interactions with male adolescents and their families regarding sexual/reproductive health requires an individualized patient-centered approach; and

3. offer health care providers specific advice on how to deliver sexual/reproductive health care to male adolescents using each encounter with a young man as an opportunity.

I. CORE MALE ADOLESCENT SEXUAL/REPRODUCTIVE HEALTH KNOWLEDGE AREAS

A. Puberty

With pubertal changes and the development of reproductive capacity come questions and concerns. Puberty for male adolescents follows a predictable sequence, but clinicians need to be aware that its timing is variable because of a variety of factors, including heredity and race/ethnicity. For boys, the first visible sign of puberty and the hallmark of the second sexual maturity rating (SMR) stage (SMR2) is testicular enlargement, followed by penile growth (the hallmark of SMR3). During SMR4, a male’s testicular volume has reached approximately 9 to 10 cm³ and his peak height growth typically occurs.

Results from a recent national survey suggest that the median age of boys’ pubertal initiation is trending earlier in the United States, especially among non-Hispanic white boys.^{14,15} One study used estimates from a sample of non-Hispanic white, non-Hispanic black, and Mexican American males 8 to 19 years of age. Two independent analyses of the same data indicate that the median age of SMR2 for genital development was 10.0 years for non-Hispanic white boys, 9.2 to 9.5 years for non-Hispanic black boys, and 10.3 to 10.4 years for Mexican American boys, whereas the median age of SMR2 for pubic hair development was 12.0, 11.2, and 12.3 years, respectively.^{16,17} These data suggest that a significant portion of boys are entering puberty earlier than 9.5 years, which was the previous lowest age of normal entry into puberty for

boys. Future studies with comparative measures are needed to confirm these secular trends.

B. Alterations in Growth Associated With Puberty

Health conditions related to male growth and development are not uncommon, can be quite distressing, and might not be identified until adolescence. The frequencies of more common disorders are 1 in 500 to 700 for Klinefelter syndrome; 1 in 1000 to 4000 for fragile X syndrome; 1 in 5000 to 10 000 for Marfan syndrome; and 1 in 8000 to 10 000 for Kallman syndrome.^{18–21} Other non-STI-related male genital issues that occur during adolescence include gynecomastia (40%–65% of male teenagers), testicular torsion (8.6 per 100 000 males 10–19 years of age), varicocele (10% of males), and testicular cancer (3.1 in 100 000 males 15–19 years of age).^{22–27}

Early and delayed pubertal timing, including short stature, can result in negative consequences for the developing male. Consequences can include higher mean levels of aggression and delinquency. Earlier-maturing boys might have more frequent involvement in risk-taking behaviors, and later-maturing boys might have lower levels of confidence and self-efficacy and increased experiences of teasing, bullying, mental health issues, and substance abuse.^{28–30} Even a common issue such as acne, which affects 95% of male adolescents,³¹ can be related to self-reported embarrassment, lower self-esteem, depression, and anxiety.³²

Given the frequency of these growth and developmental concerns, the typical practicing health care provider will certainly see male patients with 1 or more of these disorders.

C. Sexuality

Sexuality, as defined by the World Health Organization in 2002, is a cen-

tral aspect of the human life course and encompasses sex, gender identities and roles, sexual orientation, intimacy, and reproduction.¹² Although sexuality can be experienced and expressed in thoughts, fantasies, desires, beliefs, values, behaviors, roles, and relationships, not all sexuality dimensions are experienced or expressed. One's sexuality is also influenced by a variety of factors including biological, psychological, familial, societal, political, cultural, and religious factors. Before adolescence even begins, boys might be curious and ask questions about sex, body parts, differences between boys and girls, and where babies come from. However, not all parents talk about sex with their children. In the United States, many more mothers than fathers talk about sex with their sons, but mothers are less comfortable talking about sex with their sons than with their daughters.^{33,34} Fathers have also been shown to have more difficulty communicating about sexuality-related topics with their adolescents.³⁵

A recent study of California teenagers and their parents found that adolescents who reported that their parents had more frequent sexual-related communication also reported feeling closer to their parents, being more comfortable talking with their parents in general, and having more open conversations about sex with their parents.³⁶ Health care providers are in a position to advise male patients and their parents about the value and importance of repeatedly talking about sex and related topics.³⁷ Involving parents and sons in sexual health discussions can help reinforce family values and create opportunities for sons to ask clarifying questions instead of relying on misinformation provided by their peers. Male adolescents should be encouraged to talk with their health care

provider about general health and, in particular, sex, relationships, and prevention of STIs/HIV and pregnancy.^{38,39} Male adolescents cite their mothers, doctors, and nurses as their principle resources for general health care concerns³⁹ and cite doctors and other health care providers as 1 of their top 4 sources of sexual health information inclusive of parents, health classes, and television.³⁸ Schools, media, and the Internet can also play important roles in the provision of sexual health information. For example, male teenagers who received instruction on AIDS prevention and sex education at school were more likely to have fewer sexual partners, engage in fewer sex acts, and use condoms more consistently than those who did not receive such information.⁴⁰ However, adolescents should understand the potential negative influence of media and the Internet, because negative sexual health messages from these sources can lead to risky sexual behaviors.^{41,42}

D. Sexual Development

During adolescence, teenagers begin the process of developing a sexual self-concept, which involves the combination of physical sexual maturation, age-appropriate sexual behaviors, and formation of a positive sexual identity and sense of well-being. In early adolescence, boys might become preoccupied with body changes, become interested in sexual anatomy and sex, compare changes in their body with others, and explore touching and mutual masturbation. Along with the experience of spontaneous erections, ejaculation related to masturbation, and the onset of nocturnal ejaculatory events during sleep (ie, "wet dreams"), there are many reasons why preadolescent and older boys might have questions and anxieties about their emerging sexuality. Later, male adolescents begin to test their ability to attract others through dating and sex-

ual behavior. It is not uncommon for a male to have anxieties and questions about genital size and function, especially when comparing himself to others and after initiating sexual behavior.⁴³

Some male adolescents might be exploring their sexual attraction to others, and others might already recognize that they are heterosexual, homosexual, bisexual, or transgendered. Studies have found that gay male youth who have a supportive environment during the disclosure process can have a more positive psychological adjustment.^{44,45} As discussed in the American Academy of Pediatrics (AAP) statement on sexual orientation and adolescents,⁴⁶ boys who are questioning and gay and do not have supportive environments are at increased risk of social isolation, school failure, family conflict, substance abuse, depression, suicide, and stigmatization.⁴⁷⁻⁵⁰ Most sexual-minority youth are quite resilient and work through adolescent development issues. Being gay or bisexual or questioning their orientation is not in itself a problem but is a risk factor for exploring other associated risks.⁵¹ Questioning, bisexual, and gay male youth who fear being "found out" may choose not to openly discuss their concerns with potentially helpful adults such as their health care provider. Because sexual identities might be "fluid" or subject to change during adolescence, providers may find it more beneficial to inquire about sexual attractions and actual behaviors of their adolescent patients rather than asking about sexual identity (ie, how a person identifies his or her own sexuality). The health care provider might ask a series of questions such as, "A question or even a concern that a teen might have is, am I attracted to both guys and girls, just guys, or just girls. . . has that been a question for you? If so, have you been

able to answer that yet?" "Have you ever had sex with women, men, or both?" "Have you ever engaged in vaginal, oral, or anal sex?"

E. Masturbation and Spermarche

On average, the age of first male masturbation occurs between 12 and 14 years of age; most boys learn about masturbation through self-discovery. Masturbation among males is common and ranges from 36% reporting masturbating 3 to 4 times per month to 10% reporting masturbating every other day or daily.⁵² There is no evidence that masturbation is harmful in general or to one's sexual development or later adult sexual adjustment.^{53,54} However, myths related to negative consequences of masturbation persist and can result in inappropriate anxiety and/or guilt.

Before puberty, boys might masturbate to orgasm; however, no ejaculation will occur until pubertal changes commence.⁵⁵ Sperm in the ejaculate, or spermarche, typically appears during SMR3, approximately 12 to 18 months after the testes begin to enlarge. Although mature sperm production begins after the first ejaculate, a young man should be considered fertile from the time of his first ejaculation.

Health care providers can reassure male adolescents that self-masturbation is a normal behavior and can be a positive expression of sexuality and a way to delay having sex and its associated risks. Health care providers can also assist male adolescents with information and resources about normal sexual physiology and function that might not otherwise be available at home or school.

F. Sexual Behavior and Its Consequences

Sexual behavior is a normal part of development. According to the Youth Risk Behavior Surveillance, more than half (59.6%) of school-aged male teenagers

report that they have had sex by the 12th grade.² Data from the 2002 National Survey of Family Growth (NSFG) indicate that among 15- to 19-year-old male adolescents who reported never having had vaginal sex, nearly one-quarter reported engaging in oral sex or mutual masturbation behaviors.^{1,56} More recent data from the NSFG indicate that approximately one-eighth of 15- to 19-year-old males report having oral sex or other sexual contact but not vaginal sex.⁵⁷ Oral sex, in particular, is part of the repertoire of sexual behaviors and might predict involvement in other sexual behaviors.^{58–60} Thus, at a minimum, three-quarters of US male youth are reporting involvement in varying types of sexual behavior. Male adolescents also report an earlier age of sexual debut² and more sexual partners than female adolescents,^{1,2,61} which are factors known to be associated with STI-acquisition risk.^{62,63} Sexual experience increases as teenagers get older; 59.6% of 12th-grade boys have report having had sexual intercourse, compared with 33.6% of 9th-grade boys.²

A substantial number of young men also report engaging in higher-risk sexual behaviors. Data from the 2009 Youth Risk Behavior Surveillance indicate that among male high school students, 25.9% reported using alcohol or drugs before last sex, 16.2% reported 4 or more lifetime partners, and 8.4% reported initiating sex at 13 years or younger.² Data from the 2002 NSFG also showed that among 15- to 19-year-old males, 28.6% reported no condom use at last vaginal sex, 11.1% reported engaging in anal sex with a female partner, and 5.6% reported having sex with a prostitute or an HIV-infected person or often/always being high during sex.⁷

Patterns of male adolescent sexual behaviors also vary according to race/ethnicity.² Higher proportions of non-

Hispanic black high school boys (72.1%) report being sexually experienced compared with Hispanic (52.8%) or non-Hispanic white (39.6%) high school boys. Higher proportions of non-Hispanic black boys (24.9%) initiate sex at 13 years of age or younger compared with Hispanic (9.8%) and non-Hispanic white (4.4%) boys.

National US data on same-sex sexual attraction and behavior is limited. From the 1995 National Longitudinal Survey on Adolescent Health, the mean report of same-sex attraction or relationship among male adolescents was 8.4% (95% confidence interval [CI]: 7.5–9.3).⁴⁹ In the 2006–2008 NSFG, 1.8% of 18- to 19-year-olds and 2.3% of 20- to 24-year-olds reported equally to both or mostly/only same-sex attraction.⁵⁷ In the 2002 NSFG, 4.5% of male teenagers 15 to 19 years of age and 5.5% of young adult men 20 to 24 years of age reported a same-sex partner.⁶¹ In the 2006–2008 NSFG, 2.5% of male teenagers 15 to 19 years of age and 5.6% of young adult men 20 to 24 years of age reported a same-sex partner.⁵⁷ Discordance between sexual attraction/orientation and behavior is also possible, because one's sexual attraction and identity do not always predict sexual behavior.^{57,64,65}

G. Unwanted Sex

Although males are more likely than females to agree with the statement, "sex is something that just happens,"⁶⁶ the majority (82%) of males 12 to 19 years of age reported feeling pressured by friends to have sex.⁶⁷ Among sexually experienced males 15 to 19 years of age, more than half (55%) wished that they had waited longer before having sex for the first time,⁶⁸ and more than one-third (38%) of men 18 to 24 years of age reported that they really did not want sex to happen the first time that it did or had mixed feelings about it.⁵⁶ Approximately 1 in 12

men (7.6%), particularly those whose first sexual intercourse was at younger than 15 years and non-Hispanic black men, reported that they had actually been coerced to have sex by a female (5.8%) or male (2%).⁵⁶ Long-term consequences of sexual abuse of males include HIV risk behaviors, psychiatric disorders, substance abuse, and thoughts of suicide.^{69–72}

Many male adolescents and young adults are unaware of state laws on statutory rape and resultant consequences. Although these punitive measures attempt to force people to change their sexual/reproductive health behavior, there are few data to support that such laws are effective deterrents to, for example, adolescent pregnancy.⁷³ A recently prepared report by the Lewin Group contains a guide to state laws on statutory rape and reporting requirements.⁷⁴ It is recommended that health care providers be aware of local statutory rape laws and be able to counsel teenagers about potential legal consequences of early sexuality, especially if a patient's sexual partner(s) is younger than the patient. As discussed in the AAP statement on media education,⁷⁵ pediatricians also need to become educated about the public health risks of media such as "sexting." Resources for patients and families that explain more generally the laws about statutory rape, age of consent, sexting, etc (www.sexlaws.org) are also available.

H. Dating Violence

Violence in adolescent relationships can include bullying, threatening, sexual harassment, dating violence, and/or coercion. Within the context of intimate relationships among romantic and sexual partners, such violence can be verbal, emotional, physical, or sexual. Male adolescents can be perpetrators, victims, or both. For example, 9.9% of high school students na-

tionwide reported having been hit, slapped, or physically hurt intentionally by their boyfriend or girlfriend in the past year.⁷⁶ Overall, the prevalence of experiencing dating violence was higher among male (11.0%) than female (8.8%) 9th- and 12th-grade students. Despite the high prevalence, many adolescents who are victims or perpetrators of violence do not seek help.⁷⁷ Health care providers have an opportunity to promote healthy relationships, improve communication, and improve the detection of unhealthy relationships through screening for intimate partner violence (1 example of a mnemonic screening tool for violent behavior is FISTS [fighting, injuries, sex, threats, and self-defense]) and subsequent referral when appropriate.^{78–81}

I. Sexual Function and Dysfunction

Healthy sexual function has an important role in adolescents' and young adults' well-being and development. Few studies, however, have examined sexual health and/or problems with sexual dysfunction among adolescents. Common causes of sexual dysfunction among young adult men include anxiety about performance, premature ejaculation, worries about attractiveness during sex, decreased pleasure associated with condom use, and organic performance-related issues attributable to comorbid medical conditions (eg, diabetes, cardiac disease, neurologic deficits) or adverse effects from medication (eg, selective serotonin-reuptake inhibitors or alcohol).^{82,83} Health care providers have an opportunity to screen for problems with sexual performance (eg, "Some men may be worried because they ejaculate too soon or late or have difficulty getting or losing an erection. Have you ever experienced any of these problems?") and offer reassurance or appropriately assess for an underlying medical condition.

J. STIs/HIV, Pregnancy, and Adolescent Fatherhood

Involvement in sexual behaviors places young men and their partners at risk of negative sexual/reproductive health outcomes that are preventable, such as STIs, HIV infection, unintended pregnancy, and reproductive health-related cancers such as anal cancer attributable to human papillomavirus (HPV).

Although adolescents and young adults 15 to 24 years of age represent only one-quarter of the sexually active US population, this population is estimated to account for half of the 18.9 million new STI cases⁸⁴ and approximately 30% of all new HIV infections⁸⁵ that occur each year. Among STIs, chlamydia, HPV infection, and trichomoniasis account for 88% of all infections.⁸⁴ Among all populations, rates of chlamydia and gonorrhea are highest among people 15 to 24 years of age.^{86,87} The chlamydia prevalence rate for males 14 to 19 years of age reported from the recent National Health and Nutrition Examination Survey (1999–2002) was found to be 2.3% (95% CI: 1.5–3.5).⁸⁸ This rate is not too dissimilar from the chlamydia rate of 3.67% (95% CI: 2.93–4.58) reported among young adult men 18 to 26 years of age in the National Longitudinal Study on Adolescent Health. The Add Health study also reported prevalence rates among 18- to 26-year-olds for gonorrhea (0.44% [95% CI: 0.26–0.77]) and trichomoniasis (1.7% [95% CI: 1.3–2.2%]).^{87,89} It is common for males with chlamydia and gonorrhea infections to be asymptomatic. Males who are not circumcised are also more likely to have gonorrhea (adjusted odds ratio: 4.0 [95% CI: 1.9–8.4]) and syphilis (adjusted odds ratio: 1.6 [95% CI: 1.2–2.2]) infections than circumcised males.⁹⁰ Racial disparities have been observed for STIs other than HIV. Non-Hispanic black male adolescents have higher

rates of chlamydia, gonorrhea, and trichomoniasis compared with those of other race/ethnicity groups.^{85,89,91}

In 2006, an estimated one-quarter of the 1.1 million people living with HIV infection in the United States were not aware of their HIV status. Young people 13 to 24 years of age represent an estimated 10% of undiagnosed cases each year.⁹² HIV prevalence among young men who have sex with men is estimated at 7.2% (5.6% for those 15–19 years of age, 8.6% for those 20–22 years of age). Rates are higher for non-Hispanic black, mixed/other race, and Hispanic young men.⁹³ Young men at higher risk of HIV infection include men who have sex with men; have unprotected sex (especially with multiple partners); are past or present injection-drug users; exchange sex for money or drugs; are past or present sex partners of HIV-infected or bisexual persons or injection-drug users; are being treated for an STI; have a history of blood transfusion between 1978 and 1985; and request an HIV test, because nondisclosure might be indicative of high-risk behavior.⁹⁴ Male circumcision among mostly African samples has been associated with a lower risk of HIV acquisition to the male himself and possible benefits to the female by decreasing transmission.^{95–98} The Centers for Disease Control and Prevention (CDC) is currently considering guidelines for the role of male circumcision in the prevention of HIV transmission in the United States.⁹⁹

After a reduction in teen births and pregnancies from 1991 to 2005, more recent trends in behavioral risks for pregnancy (eg, declines in contraceptive use) have stalled or are reversing among certain groups, particularly non-Hispanic black and Hispanic teenagers.¹⁰⁰ National data indicate that approximately 750 000 pregnancies occur each year among women younger than 20 years¹⁰¹; the majority of teen

pregnancies (82%) are unplanned or mistimed.¹⁰² These data translate to approximately 1 in 20 female teenagers becoming pregnant each year.¹⁰³ Most of these pregnancies end in a live birth (57%), 27% end in induced abortion, and 16% end in fetal loss.¹⁰⁴ The majority of teen pregnancies are fathered by males who are within 2 years of age or older of their partners.^{105–108} Among sexually experienced male adolescents, 1 in 8 (13%) reported that they have impregnated a partner, and 4% are fathers.¹⁰⁹ Because of lacking or inaccurate documentation of paternal age on birth certificates, especially among younger and unmarried mothers,¹¹⁰ the rates of fatherhood among male adolescents might be even higher.

The consequences of teen pregnancy and STIs, including HIV infection/AIDS, are many and costly. Males diagnosed with an STI can have complications such as epididymitis and infertility^{111,112} and, if left untreated, can place female partners at risk for morbidity and mortality, such as acute and recurrent pelvic inflammatory disease and resultant tubal scarring, infertility,^{113–115} ectopic pregnancy,¹¹⁶ and HIV infection/AIDS. Although it can be challenging to disentangle the link between teen pregnancy and poverty, personal costs of teen pregnancy often include truncated schooling, in-

creased unemployment, welfare dependency, lower lifetime earnings, single parenthood, depression, and poor infant outcomes.^{117,118} Each year, costs related to adolescent pregnancy and its complications are estimated to be \$9 billion,^{119,120} and costs related to STIs/HIV infection and their complications are estimated to be \$17 billion.⁸⁶

K. Preventing STIs/HIV Infection and Pregnancy

The male adolescent and his partner, family, school, and health care provider all have important roles in preventing the negative consequences of sexual behavior.

Table 1 summarizes the effectiveness of available male contraceptive options.¹²¹ Condoms are the most commonly used contraception method¹²² and are highly effective if used consistently and correctly. Over the past 3 decades, the report of condom use by male adolescents at first and most recent sex has increased, particularly among Hispanic and non-Hispanic black males.¹ Seven in 10 male adolescents (71%) report condom use the first and most recent time they had sex.¹ The majority of young men also report using condoms to prevent both pregnancy and STIs/HIV infection.⁵⁶ However, during the past 3 decades, there has been little change in male

TABLE 1 Contraceptive Effectiveness: Rates of Unintended Pregnancies per 100 Women

	First-Year Pregnancy Rate, % ^a		Key
	Consistent and Correct Use	As Commonly Used	
Male methods			
Vasectomy	0.10	0.15	0.0%–0.9%: very effective
Male condom	2	15	1%–9%: effective
Withdrawal	4	27	10%–25%: moderately effective
No method	85	85	26%–32%: less effective
Common female hormonal methods			
Progestin-only injectable	0.3	3	—
Combined oral pills, patch, ring	0.3	8	—

^a Rates are largely from the United States.

Modified from: Trussell J. Contraceptive efficacy. In: Hatcher RA, Trussell J, Nelson AL, et al, eds. *Contraceptive Technology*. 19th Rev ed. New York, NY: Ardent Media; 2007:747–826.

adolescents' consistency of condom use (ie, use with every sexual encounter in the past year); less than half (48%) of sexually active males report consistent use.¹ Barriers to condom use include being embarrassed when buying them, reduced physical sensation, incorrect use and lack of availability, planning, discussion, or negotiation with one's partner.^{1,123–128} Reduced likelihood of condom use at most recent sex and inconsistency in condom use have been observed among young men who did not receive formal sex education at school or the community.¹²⁹ Low condom use during adolescence, combined with the report of multiple partners, has also been shown to be associated with elevated STI rates when subjects become young adult men.¹³⁰

Approximately 10% of male adolescents report using withdrawal as their contraception method of choice.¹ Although withdrawal does not protect against STIs/HIV infection and, thus, is not an appropriate form of STI/HIV prevention, effectiveness data (Table 1) show that it can substantially reduce the risk of pregnancy for people who report difficulties using contraception (eg, people with latex allergy or during unplanned sex acts).¹³¹ Withdrawal is better than doing nothing, but it must be emphasized that withdrawal is not equivalent to using other barrier methods (eg, condoms), hormonal methods, or dual contraception. Polyurethane condoms are also available for people who are allergic to latex.

Only one-quarter (24%) of male adolescents report the use of dual contraception methods (eg, combined condom use with a hormonal method).¹ This may be explained by a lack of awareness of the partner's contraception methods or familiarity with how female hormonal methods work, including emergency contraception.^{132–134} For ex-

ample, situations in which a male was less likely to use a condom included if the male's partner used a method of contraception, if the male was older at most recent sex, if the male's partner was older at first sexual encounter, and during a casual first sexual encounter.¹²⁹ Increased odds of ever or always using any contraceptive method is observed among teenagers who wait longer between the start of a relationship and first sex, discuss contraception before first sex, or use dual contraception methods.¹³⁵ Two groups of teenagers, males and non-Hispanic white teenagers, are less likely to regularly talk about contraception and STIs with their partners before first sex than are females and teenagers from other racial groups.¹³⁶ Health care providers have an opportunity to talk with males about all types of contraception and provide consistent messages about counseling for risk reduction (Table 2). Some teenagers might choose to take virginity pledges and abstain from sex until marriage. However, such pledges have not been found to have long-term effectiveness in preventing STIs and other adverse outcomes and might actually place pledge-takers at greater risk of STIs and unintended pregnancy. One prospective study found that adolescents who signed abstinence pledges experienced similar STI rates as did adolescents who did not sign pledges.¹³⁷ Although pledge-takers delayed initiation of sex for a longer period of time, when pledge-takers initiated sex, they were less likely to use condoms and seek reproductive health care compared with their peers who did not sign pledges.

L. Male Adolescents' Sexual/Reproductive Health Care Use

Male adolescents' use of sexual/reproductive health care services remains low for a variety of reasons. For many, the onset of sexual behavior during ad-

olescence serves as a rite of passage into manhood but does not trigger thinking about preventive behaviors related to the consequences of sexual behavior. Young men might also hold more traditional masculine beliefs that preclude them from seeking care despite having symptoms.^{138,139} In addition, the components of young men's sexual/reproductive health care have been poorly defined and have historically received little attention. Unlike females, who have historically received bundled sexual/reproductive health care as part of gynecologic examinations, birth control visits, and prenatal care,¹⁴⁰ this is not typically true for young men. Barriers to care can also play a role in male adolescents' care-seeking in general and sexual/reproductive health care in particular. These barriers include fear; stigma; shame; denial; lack of social support; lack of confidential services; lack of health insurance options especially as they get older; and not knowing where to go for care.^{141–146} Given the asymptomatic nature of many STIs, including HIV infection, strategies are needed to raise young men's awareness about sexual/reproductive health services, STI and HIV testing options, and related resources. Promoting sexual/reproductive health for young men can lead to sexual responsibility, healthy intimate relationships, and responsible fatherhood. Health care providers, and pediatricians in particular, are in the best position to deliver high-quality sexual/reproductive health care services to male adolescents and should view even follow-up, acute care, and immunization visits as opportunities to address these health issues. Health care providers can also use "clinical hooks," such as sports physicals or acne follow-up, to keep male adolescents engaged in care and to deliver sexual/reproductive health care services.

TABLE 2 Anticipatory Guidance Topics

Anticipatory Guidance Topics and Questions	Example Message	Resources
Counseling on sexual risk reduction including barriers to condom use 2 separate sessions, 1 wk apart	Avoiding sex is the safest way to prevent pregnancy and STIs/HIV. If you choose to have sex, take responsibility and use a condom <i>everytime</i> . If you do not have a condom, choose not to have sex.	www.ahwg.net (see sexual health toolkit—provider, pages C13–C15)
Session 1: assess for personal risk, barriers to risk reduction, and identification of a small risk-reduction step within 1 wk (eg, consistent condom use)	Choosing not to have sex at any point will not diminish one's status or manhood.	www.sexetc.org/topic/guys_health
Session 2: review previous week's behavioral-change successes and barriers, provide support for changes made, identify barriers and facilitators to change, and develop a long-term plan for risk reduction		www.goaskalice.columbia.edu/Cat7.html
Help me understand why condoms are difficult for you to use. Do you vary your condom use depending on who your partner is? How can you reintroduce condom use in a relationship without creating issues about trust?		www.iwannaknow.org
What would you do if your partner does not want you to use a condom?	"I really like you but am not comfortable having sex without using a condom."	
Do you know how to make condom use more pleasurable?	Place a drop of lubrication in the inside tip of the condom. Also, condoms come in various shapes and sizes, and 1 condom type may not fit all.	www.ahwg.net (see sexual health toolkit—provider, pages C36–C37; youth page C50)
What would you do if a condom breaks or slips off?	"Do you know about emergency contraception (eg, Plan B)? This is something we can get and have you take within the next 3 to 5 days to help prevent you from getting pregnant if you are not on birth control."	www.not-2-late.com ; 1-888-NOT-2LATE; http://bedsider.org/where_to_get_it
What do you know about the consequences of sexual behavior?	You can have an STI and not even know it. Not all STIs have symptoms, and not all STIs are curable. Getting pregnant as a teenager comes with many legal and economic responsibilities. Having sex can have legal implications depending on who you have sex with and how old you and your partner are.	www.youngmenshealthsite.org/std-general.html www.babycenter.com/babyCostCalculator.htm www.sexlaws.org
Demonstrate or provide resources (eg, videos, handouts) for correct condom use; distribute condoms		
Have you ever been taught, using a condom model, how to correctly put on a condom?	Using a condom can help reduce your risk of getting an STI and/or getting someone pregnant.	www.cdc.gov/condomeffectiveness/brief.html ; www.stdcentral.org/SitC/about
How comfortable are you with putting on a condom?	Check that the condom has not expired and has been properly stored. When opening the package, be careful not to rip the condom. Once open, place the correct side of the condom over an erect penis. Hold onto the reservoir tip at the top, remove any excess air and slowly roll the condom all the way down over the erect penis to the penis base. After ejaculation, hold onto the condom rim at the base of the penis, making sure the condom does not slip off or leak. After removing the condom, dispose of it in the trash.	
Communication with partner about sex		
Do you talk with your partner about sex, preventing STIs/HIV, getting tested for STIs/HIV before having sex for the first time, getting tested together, preventing pregnancy, whether she is on birth control?	I encourage you to talk with your partner about contraception and getting tested together before you start engaging in a new sexual relationship.	www.ahwg.net (see sexual health toolkit—provider pages C45–C65)
Do you know about the different types of female birth control methods?		www.youngwomenshealth.org/contratypes.html

TABLE 2 Continued

Anticipatory Guidance Topics and Questions	Example Message	Resources
<p>Help teenaged fathers understand their role in parenting</p> <p>Help me understand your relationship with your child(ren) (eg, emotional, physical, and financial support and barriers/facilitators to involvement)?</p> <p>Assess for experience of stress</p> <p>Identify local community resources to promote parenting role (parenting classes, educational support, job training, etc)</p>	<p>We know that fathers who are involved in their children's lives can improve the health of their children and the health of the fathers themselves.</p>	<p>www.fatherhood.org</p>

II. ROLE OF THE HEALTH CARE PROVIDER

A. Provide Individualized Sexual/Reproductive Health Care to Male Adolescents

The provision of sexual/reproductive health care to young men in the clinical setting should be individualized on the basis of the patient's developmental and psychosocial needs by using patient-centered approaches.¹⁴⁷ Health care providers need to be comfortable taking a young man's sexual/reproductive history to elicit the type of information needed to determine the appropriate anticipatory guidance, counseling, and treatment. Trust and relationship-building are also critical elements of the male adolescent's visit that help him to feel comfortable regardless of a physician's gender and/or background. Interviewing the young man one-on-one, separate from accompanying parents or guardians, will also improve the quality of information received and ensure confidentiality of the information shared. As with young women, health care providers need to be cognizant of minor consent laws in the states in which they practice and educate patients about specific state laws as appropriate.^{148,149}

Myths about adolescent male sexuality still pervade American society. Health care providers should avoid making assumptions or allowing stereotypes about male adolescent sexuality influence interactions with their young

male patients. Health care providers need to understand the specific needs of gay, bisexual, or questioning male adolescents and provide support and/or referral resources as appropriate. Sexual initiation during adolescence should be viewed as part of sexual development. Contrary to the image of male promiscuity, the majority of male adolescents have no more than 1 sex partner during a 1-year period. Most male adolescents also believe that they should be responsible for pregnancy prevention and report condom use as a means of preventing both STIs and pregnancy. Rather than holding young men responsible for the negative sexual/reproductive health outcomes seen among young women (pregnancy, STIs/HIV, and relationship violence), health care providers can address the other half of the partner equation that has historically been left out. Health care providers should share clear and consistent messages with all male adolescents, their parents, and schools about sexual health, comprehensive sex education, and regular care-seeking.

B. Deliver High-Quality Sexual/Reproductive Health Care to Male Adolescents

A number of national organizations make recommendations about clinical services that health care providers should deliver, including the AAP's *Bright Futures*,⁴ the CDC,¹⁵⁰⁻¹⁵² and the US Preventive Services Task Force (USPSTF).¹⁵³ Sexual/reproduc-

tive health services that have been shown to be most effective for health care providers to provide include taking sexual health histories, assessing sexual risk, and providing behavioral counseling regarding STIs.^{4,150,153} Regarding laboratory screening for HIV infection, the USPSTF recommends HIV testing on the basis of risk factors for the individual and/or clinic setting.^{150,154} The CDC and *Bright Futures* recommend HIV testing for all persons 13 years of age and older and subsequent testing for all persons at high risk at least annually.^{4,150,154} Regarding laboratory screening for STIs, although the USPSTF has stated that there is insufficient evidence to screen sexually active young men for chlamydia and gonorrhea,^{155,156} the CDC recommends chlamydia screening among sexually active young men in high-prevalence settings (eg, STI, Job Corps, or jail clinics) and rescreening of persons infected with chlamydia 3 months later,^{151,152} and *Bright Futures* recommends screening sexually active adolescents for chlamydia and gonorrhea.⁴ The USPSTF¹⁵⁷ and *Bright Futures*⁴ also recommend testing for syphilis on the basis of individual risk factors. *Bright Futures* also recommends an annual genital examination for SMR and observations for signs of STIs (ie, warts, vesicles) and testicles for hydrocele, hernias, varicocele, or masses. Refer to the AAP's recommended childhood and adolescent immunization schedule.¹⁵⁸ The CDC Advi-

sory Committee on Immunization Practices' recent update recommends routine vaccination of male adolescents against HPV starting at age 11 through 12 years, states that the series can be started beginning at age 9 years, and recommends catch-up vaccination among males ages 13 through 21 years who have not been vaccinated previously or have not completed the 3-dose series through age 21 years. Such vaccination is also recommended in AAP immunization schedule (for updates, see: www.aap.org/immunization/izschedule.html).^{158,159} Males age 22 through 26 years may be vaccinated (permissive recommendation for this age group). Routine vaccination is recommended among at-risk males including males who have sex with males and immunocompromised males through age 26 years.

The delivery of sexual/reproductive health services is one of the few effective clinical preventive services that can positively affect male adolescents' and young adults' health.¹⁶⁰ However, health care providers miss many opportunities to assess young men's sexual health and deliver STI/HIV services; on average, fewer than one-quarter of male adolescents and young adult men report receiving sexual/reproductive health services compared with more than half of similarly aged female adolescents.^{9,10,161,162}

Health care providers can use each encounter in the office setting as an opportunity to address the core recommended components of male adolescents' sexual/reproductive health with all male adolescent patients, including the provision of appropriate resources to patients and parents. Health care providers should consider scheduling an annual sexual/reproductive health visit for the sexually active young man. The health care provider should know how to:

- take a complete sexual history;

- perform a complete sexual/reproductive health examination that includes examination of the skin, breasts, genitals, hair, and, if relevant, the perianal area and documentation of the SMR;
- perform and interpret STI/HIV tests and know how to treat STIs;
- administer vaccinations, as recommended by the AAP,¹⁵⁸ to protect against infections that are transmitted sexually, such as hepatitis B and A vaccines and HPV vaccine¹⁵⁹;
- provide anticipatory guidance/counseling on sexual/reproductive health matters (Table 2), including the use of messages about dual methods (eg, "not having sex is the best way to avoid pregnancy and STIs/HIV, but if you choose to have sex, use condoms consistently and use a reliable contraceptive method for the partner"); and
- use resources (Table 3) in varying formats (ie, clinic handouts, books, Web-based resources, community resources) to share and/or refer patients and parents as needed.

C. Take a Sexual History

A comprehensive and confidential sexual assessment is a standard part of the past medical, family, and social history and should begin during early adolescence and be updated at least annually. This can be accomplished face-to-face or through the use of confidential pre-visit questionnaires (Table 3).^{163–165} Components of a comprehensive sexual health assessment should include asking about attractions to females, males, or both; sexual initiation and age of onset; engagement in vaginal, oral, or anal sex; gender of partner(s); number of partners; current STI symptoms or concerns; history and results of STI/HIV testing; use of condoms; partners' use of methods to protect against STIs and/or pregnancy; and ex-

perience of sexual abuse. Assess for comfort with changes in one's body, masturbation, knowledge about how to use a condom correctly, hormonal contraception methods, and emergency contraception. Among sexually experienced males, assess pressure to have sex, access to condoms, use of alcohol or other substances when having sex, experience of pregnancy or fatherhood, experience of being a perpetrator or victim of verbal/physical abuse or force to have sex in a relationship, whether sex is pleasurable, and/or problems with sexual performance. Among patients who state a high number of partners or are hesitant to share their number of partners, consider and ask about whether they are exchanging sex for money or drugs. Among non-sexually active young men, assess for dating, pressure or plans to initiate sex, and knowledge about ways of preventing STIs/HIV and pregnancy. During childhood, encourage and provide resources to parents and families to have developmentally appropriate conversations with their sons that continue through adolescence about sexuality, sex, and other sensitive topics.³⁷

A number of verbal and written tools are available that can assist health care providers to take a sexual health assessment and other components of the adolescent's psychosocial history,¹⁶⁶ including the HEEDSSS (home, education and employment, eating, activities, drugs, sexuality, suicide and depression, and safety) assessment and *Bright Futures* toolkits (http://brightfutures.aap.org/tool_and_resource_kit.html).^{4,167,168}

D. Perform a Physical Examination

Despite the lack of evidence-based guidelines supporting routine testicular screening and teaching of testicular self-examination for detection of testicular cancer, a genital

TABLE 3 Male Teen Resources for Patients, Parents, and Health Care Providers

Male patient resources	
Brochures to order	
ETR Publishing	http://pub.etr.org
Office of Population Affairs Clearinghouse	www.hhs.gov/opa/pubs/download_pubs/download_pubs.html
Planned Parenthood	www.plannedparenthood.org/teen-talk
Downloadable brochures and male-friendly Web sites	
American Social Health Association	www.iwannaknow.org
Children's Hospital Boston	www.youngmenshealthsite.org
Gay and Lesbian Medical Association: Online Health Care Referrals	www.glma.org
Go Ask Alice! (Columbia University)	www.goaskalice.columbia.edu/Cat7.html
Family Acceptance Project (Marian Wright Education Institute: resource for lesbian, gay, bisexual, transgendered youth and families)	http://familyproject.sfsu.edu
Palo Alto Medical Foundation (a Sutter Health affiliate)	www.pamf.org/teen/health/malehealth
Sex Etc (Rutgers University)	www.sexetc.org/topic/guys_health
TeensHealth (Nemours Foundation)	www.kidshealth.org/kid/grow
	www.kidshealth.org/teen/sexual_health
The Trevor Project	www.thetrevorproject.org/lifelinechat
Youth Resource (advocates for youth)	www.amplifyyourvoice.org/youthresource
Parent resources	
American Academy of Child and Adolescent Psychiatry (handouts on talking to your kids about sex; gay, lesbian, and bisexual adolescents)	www.aacap.org/cs/root/facts_for_families/facts_for_families
AAP Section on Adolescent Health	www.aap.org/moc/AdolHandouts_AAPMbrs/Resources.htm
Madaras L, Madaras A. <i>My Body, My Self: For Boys</i> . New York, NY: Newmarket Press; 2007	
McCoy K, Wibbelsman C. <i>The Teenage Body Book</i> . New York, NY: Perigee; 2008	
The National Campaign to Prevent Teen Pregnancy	www.thenationalcampaign.org/parents/default.aspx
Parents, Families and Friends of Lesbians and Gays	www.pflag.org
Provider resources	
AAP Section on Adolescent Health	www.aap.org/moc/AdolHandouts_AAPMbrs/ProviderHandouts.htm
Adolescent Health Working Group: an adolescent provider toolkit series on sexual health, behavioral health, body basics, and adolescent health care 101	www.ahwg.net (see toolkit provider resource page)
American Medical Association Guidelines for Adolescent Preventive Services (AMA GAPS): previsit questionnaires	www.ama-assn.org/ama/pub/physician-resources/public-health/promoting-healthy-lifestyles/adolescent-health.shtml
<i>Bright Futures</i> : toolkit	http://brightfutures.aap.org/tool_and_resource_kit.html
CDC: expedited partner therapy and places for referral to local public health facilities for STI/HIV testing	www.cdc.gov/std/ept ; www.hivtest.org
Physicians for Responsible Choice and Health	www.prch.org
Society for Adolescent Health and Medicine: previsit questionnaires	www.adolescenthealth.org/Clinical_Care_Resources/2173.htm

examination, including examination of the testicles, represents an important part of a male adolescent's complete physical examination during annual preventive health visits and, specifically, as part of a visit related to a genital complaint.⁴ The content of the pediatric/adolescent physical examination required to report preventive health care codes (*Current Procedural Terminology* [CPT] codes 99382–99384; 99392–99394) depends on age and developmental

level and would be expected to include a male genital examination. There are many reasons to perform a genital examination, including SMR for hair and genital progress in pubertal development; screening for visual signs of STIs such as herpes, warts, and asymptomatic urethral discharge; identification of signs for genetic diseases such as Klinefelter syndrome or ambiguous genitalia (congenital adrenal hyperplasia); evidence of structural anomalies

including varicocele, which can affect fertility, or uncorrected hypospadias, which can result in significant embarrassment, problems with sexual function, and/or abnormal urine flow. Other issues might be related to an uncircumcised penis (eg, phimosis and hygiene); hair or skin findings for treatable conditions (eg, folliculitis and tinea cruris); previously unrecognized absent testes attributable to cryptorchidism; evidence of testicular atrophy secondary to central or exog-

enous causes (eg, steroids or marijuana use); and normal findings that might require reassurance (eg, pearly penile papules).^{111,168,169}

Ultimately, one of the goals of the genital examination is to help the young man gain a better understanding of his own body and reproductive parts. For example, young men might not be aware of their genital anatomy, the importance of using genital protection as part of sports participation, or general issues related to their hygiene. Guidance can be given by briefly reviewing the main components of the physical examination and, as the examination progresses, by commenting on normal anatomy, what to expect next, and pertinent findings. Examination in gowns will help prevent missing important physical examination findings, such as gynecomastia or truncal acne. Avoiding lengthy discussions while the patient is undressed also allows for greater patient comfort.

Despite the lack of recommendations that support testicular cancer screening, the USPSTF has noted that “clinicians should be aware that patients who present with symptoms of testicular cancer are frequently diagnosed as having epididymitis, testicular trauma, hydrocele or other benign disorders.”^{170,171} The American Cancer Society has stated that it “does not have a recommendation on regular testicular self-examinations for all men.”¹⁷² Risk factors for testicular cancer among men include being white; being between 13 and 39 years of age; having a history of cryptorchidism, testicular atrophy/dysgenesis, testicular trauma, HIV infection, and Klinefelter syndrome; or having a family history of testicular cancer. An external anal inspection, a digital rectal examination, and screening for hernia as part of the male adolescent physical examination should be performed on the basis of specific concerns or complaints such as a

bulging mass or pain (hernia examination), hemorrhoid or rectal bleeding (digital rectal examination), or risk factors that would warrant an external anal inspection for HPV lesions in a young man who engages in receptive anal intercourse.

Health care providers might be confronted with male adolescents who refuse a genital examination because of concerns about homophobia, lack of experience with such examinations, fear of getting an erection, or even because of previous abuse. Understanding the specific concern can help the health care provider educate the patient about the importance of this examination, determine the priority of such an examination for a particular patient, and negotiate how and when to complete the required components of the examination. Routinely examining the genitals from childhood through adolescence can help the male patient understand the routine nature of this examination component. The use of a chaperone might also be relevant and should be considered during all genital examinations for patient and/or provider comfort regardless of whether the provider and patient are the same gender.¹⁷³

E. Perform Laboratory Screening

Sexually active male adolescents 13 years of age and older should be offered an HIV test. HIV testing should occur at least once by 16 to 18 years of age or sooner once an adolescent is known to be sexually active or otherwise at risk. Youth at high risk should be tested annually for HIV infection. Adolescents tested for other STIs should be tested for HIV at the same visit. Youth seen in urgent care facilities and emergency departments should be routinely tested for HIV. In addition, routine HIV screening of young persons in high-risk or high-prevalence settings, such as juvenile

detention centers and jails, STI clinics, and teen clinics, has been shown to increase the yield for HIV screening. HIV testing options include serologic tests that provide results within 1 to 2 weeks and newer noninvasive oral rapid tests that provide results within 20 minutes.

Sexually active male adolescents are considered to be among the high-risk groups for STIs and, in general, should be screened even if they are asymptomatic for chlamydia using noninvasive urine tests. Other screenings available as urine tests include those for gonorrhea and trichomonas.⁸⁹ Routine screening of asymptomatic males should be based on the prevalence rates within the communities. Among sexually active male adolescents with painful urination (dysuria) and/or urethral discharge, urethritis attributable to an STI will be the most common cause of infection, as opposed to urinary tract infection, which affects less than 0.01% of male adolescents with normal urogenital anatomy.¹⁷⁴ Noninvasive tests are the preferred method (eg, urine-based nucleic acid amplification tests [NAATs] to screen for chlamydia and gonorrhea). These tests have high sensitivities and specificities (>90%) as well as high patient acceptability. Refer to the current CDC's sexually transmitted diseases treatment guidelines for best testing strategies if NAATs are unavailable.¹⁵²

For male patients who report sexual behavior with another man, the CDC's sexually transmitted diseases treatment guidelines recommend HIV serologic testing, if HIV-negative or not tested in the previous year; syphilis serologic testing; a test for urethral infection with chlamydia and gonorrhea among men who have had insertive sex during the preceding year; a test for rectal infection with chlamydia and gonorrhea among men who have had receptive anal sex during the preceding year; and a test for pharyngeal in-

fection for gonorrhea among men who have had receptive oral sex during the preceding year.¹⁵² No current guidelines detail screening recommendations for chlamydia of the pharynx or for anal screening of the rectum using anal Papanicolaou tests or HPV typing.^{175–177} Anal Papanicolaou tests and/or HPV typing are currently under review as possible screening tools for anal carcinoma. Nucleic acid amplification tests are not currently approved by the Food and Drug Administration for pharyngeal or rectal specimens but might be available for use in some laboratories. Both chlamydia and gonorrhea can infect the rectum, and gonorrhea can also infect the pharynx. However, there is little evidence that chlamydia can cause pharyngeal infection.

Health care providers should be able to diagnose STIs that occur most commonly in male adolescents and young adults, including chlamydia, gonorrhea, HPV, trichomoniasis, herpes, ascending infections (ie, epididymitis), and syphilis. The CDC's sexually transmitted diseases treatment guidelines (www.cdc.gov/std/treatment) and the AAP's *Red Book* (<http://aapredbook.aappublications.org>) are updated on a regular basis and provide the most up-to-date screening, testing, and treatment recommendations.^{152,178} Health care providers should also encourage STI treatment of their partners through provision of expedited partner therapy, if allowed by state law¹⁷⁹ (www.cdc.gov/std/ept); bringing the partner in for testing and treatment; or referral to local public health facilities (www.hivtest.org).

Health care providers are encouraged to communicate to their patients and families that the delivery of sexual/reproductive health services, including STI/HIV testing, is a standard and routine part of adolescent clinical services. State mandates that require an

explanation of benefits to be sent to parents of adolescent patients might conflict with patients' rights to confidential care and minor consent. Adolescents who are seeking confidential testing for STIs might require referral to settings that provide confidential STI testing, such as local public health clinics or Planned Parenthood centers. Informed and proactive health care providers can ensure that their patient's confidentiality is maintained.

F. Provide Anticipatory Guidance and Counseling

Health care providers should tailor anticipatory guidance and counseling on the basis of specific risk factors and/or information obtained during an individual male patient's visit and his developmental stage. Health care providers should continue to be up-to-date on coding and local reimbursement practices that allow for the time it takes to counsel on preventive health care matters to be included within a problem-focused visit. Table 2 lists suggested sexual/reproductive health anticipatory guidance topics and messages for male adolescent patients.

The USPSTF recommends moderate- to high-intensity behavioral counseling in the clinic setting to prevent STIs for all sexually active adolescents.¹⁵³ One example of such a counseling approach would be 2 separate 20-minute clinical sessions 1 week apart.¹⁸⁰ During the first clinical session, patients are assessed for personal risk, barriers to risk reduction, and identifying a small risk-reduction step within the next week. During the second clinical session, the previous week's behavioral-change successes and barriers are reviewed, support for changes made is provided, barriers and facilitators to change are identified, and a long-term plan for risk reduction is developed. A study of this method found that participants in intervention clinics who were

receiving structured behavioral counseling reported significantly higher condom-use rates and fewer new STIs than participants at control sites. Males and females benefited equally from counseling interventions, and brief interventions were more effective among adolescents than older participants.

G. Administer Vaccinations

Reproductive health-related vaccines, including hepatitis A and B and HPV vaccines, should be administered to male adolescents as recommended by the AAP and the CDC's Advisory Committee on Immunization Practices along with other routine immunizations for male adolescents.

III. GUIDANCE FOR HEALTH CARE PROVIDERS

1. Know the core sexual/reproductive health knowledge areas for male adolescents and understand the affect that sexual health has on the male adolescent's quality of life.
2. Encourage parents to talk about pubertal development and have age-appropriate discussions about sexual health with their sons on repeated occasions.
3. Guide male adolescents to build healthy, responsible relationships.
4. Recognize that some male adolescents might be dealing with issues of sexuality that can affect their psychosocial and physical health, provide appropriate support for families,¹⁸¹ and identify local resources for referral when appropriate.
5. Regardless of reason for the visit and clinical setting, routinely provide quality male sexual/reproductive health services to all male adolescents.
6. Deliver core sexual/reproductive health services to male adoles-

cents: take a sexual history, perform an appropriate examination, screen for STIs/HIV, provide patient-centered and age-appropriate anticipatory guidance, and deliver appropriate vaccinations.

7. Provide sexual/reproductive health services in a confidential and culturally appropriate manner.
8. Use “clinical hooks,” such as sports physicals and acne follow-up visits, to keep young men enrolled in general and especially sexual/reproductive health care. Consider promoting an annual sexual/reproductive health visit for male adolescents to address their core sexual/reproductive health issues.
9. Include in clinic settings everything

from handouts to posters to make male adolescents feel welcome and comfortable. Educational materials such as handouts, pamphlets, references, and videos should also be made available to reinforce office-based educational efforts.

10. Continue to be up-to-date on coding and local reimbursement practices that allow for preventive health care counseling to be included within problem-focused visits.

LEAD AUTHORS

Arik V. Marcell, MD, MPH
Charles Wibbelsman, MD
Warren M. Seigel, MD

COMMITTEE ON ADOLESCENCE, 2010–2011

Margaret J. Blythe, MD, Chairperson

William P. Adelman, MD
Cora C. Breuner, MD, MPH
David A. Levine, MD
Arik V. Marcell, MD, MPH
Pamela J. Murray, MD
Rebecca F. O'Brien, MD

PAST MEMBERS

Paula K. Braverman, MD
Warren M. Seigel, MD
Charles Wibbelsman, MD

LIAISONS

Rachel J. Miller, MD – *American Congress of Obstetricians and Gynecologists*
Jorge L. Pinzo, MD – *Canadian Paediatric Society*
Benjamin Shain, MD – *American Academy of Child and Adolescent Psychiatry*

STAFF

Mark Del Monte, JD
Karen Smith

REFERENCES

1. Abma JC, Martinez GM, Mosher WD, Dawson BS. Teenagers in the United States: sexual activity, contraceptive use, and childbearing, 2002. *Vital Health Stat* 23. 2004;(24):1–48
2. Centers for Disease Control and Prevention. Youth Risk Behavior Surveillance: United States, 2009. *Morb Mortal Wkly Rep*. 2010;59(SS-5):1–142. Available at: www.cdc.gov/HealthyYouth/yrbs/index.htm. Accessed June 22, 2010
3. Marcell AV, Klein JD, Fischer I, Allan MJ, Kokotailo PK. Male adolescent use of health care services: where are the boys? *J Adolesc Health*. 2002;30(1):35–43
4. Hagan JF Jr, Shaw JS, Duncan P, eds. *Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents*. 3rd ed. Elk Grove Village, IL: American Academy of Pediatrics; 2008
5. American Medical Association. *Guidelines for Adolescent Preventive Services*. Chicago, IL: American Medical Association; 1994
6. Green M, eds. *Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents*. Arlington, VA: National Center for Education in Maternal and Child Health; 1994
7. Marcell AV, Bell DL, Lindberg LD, Takruri A. Prevalence of STI/HIV counseling services received by teen males, 1995 to 2002. *J Adolesc Health*. 2010;46(6):553–559
8. Ma J, Wang Y, Stafford RS. U.S. adolescents receive suboptimal preventive counseling during ambulatory care. *J Adolesc Health*. 2005;36(5):441
9. Lafferty WE, Downey L, Holan CM, et al. Provision of sexual health services to adolescent enrollees in Medicaid managed care. *Am J Public Health*. 2002;92(11):1779–1783
10. Lafferty WE, Downey L, Shields AW, Holan CM, Lind A. Adolescent enrollees in Medicaid managed care: the provision of well care and sexual health assessment. *J Adolesc Health*. 2001;28(6):497–508
11. International Conference on Population and Development. Summary of the ICPD Programme of Action. Cairo, Egypt: United Nations Department of Public Information; 1994. Available at: www.unfpa.org/public/cache/offonce/home/sitemap/icpd/International-Conference-on-Population-and-Development/ICPD-Summary. Accessed October 5, 2011
12. World Health Organization. *Defining Sexual Health: Report of a Technical Consultation on Sexual Health, 28–31 January 2002, Geneva*. Geneva, Switzerland: World Health Organization Press; 2006. Available at: www.who.int/reproductivehealth/publications/sexual_health/defining_sh/en/index.html. Accessed October 5, 2011
13. Lindberg LD, Sonfield A, Gemmill A. Reassessing adolescent male sexual and reproductive health in the United States: research and recommendations. *Am J Mens Health*. 2008;2(1):40–56
14. Euling SY, Herman-Giddens ME, Lee PA, et al. Examination of US puberty-timing data from 1940 to 1994 for secular trends: panel findings. *Pediatrics*. 2008;121(suppl 3):S172–S191
15. Herman-Giddens ME, Steffes J, Dowshen S, et al. Age of onset of selected pubertal characteristics: cross-sectional data from non-Hispanic white boys [abstr]. E-PAS. Available at: www.aap.org/research/abstracts/2010/8abstract10.pdf. Accessed February 1, 2010
16. Sun SS, Schubert CM, Chumlea WC, et al. National estimates of the timing of sexual maturation and racial differences among US children. *Pediatrics*. 2002;110(5):911–919
17. Herman-Giddens ME, Wang L, Koch G. Secondary sexual characteristics in boys: estimates from the national health and nutrition examination survey III, 1988–1994. *Arch Pediatr Adolesc Med*. 2001;155(9):1022–1028
18. Bojesen A, Juul S, Gravholt CH. Prenatal and postnatal prevalence of Klinefelter syndrome: a national registry study. *J Clin Endocrinol Metab*. 2003;88(2):622–626
19. O'Donnell WT, Warren ST. A decade of molecular studies of fragile X syndrome. *Annu Rev Neurosci*. 2002;25:315–338
20. Freed LA, Levy D, Levine RA, et al. Preva-

- lence and clinical outcome of mitral-valve prolapse. *N Engl J Med*. 1999;341(1):1–7
21. Dodé C, Hardelin JP. Kallmann syndrome. *Eur J Hum Genet*. 2009;17(2):139–146
 22. Mansbach JM, Forbes P, Peters C. Testicular torsion and risk factors for orchiectomy. *Arch Pediatr Adolesc Med*. 2005;159(12):1167–1171
 23. Horner MJ, Ries LAG, Krapcho M, et al. SEER cancer statistics review, 1975–2006. Available at: http://seer.cancer.gov/csr/1975_2006. Accessed June 22, 2010
 24. Feber KM, Kass EJ. Varicocelectomy in adolescent boys: long-term experience with the Palomo procedure. *J Urol*. 2008;180(4 suppl):1657–1659; discussion 1659–1660
 25. Lanitis S, Starren E, Read J, et al. Surgical management of gynecomastia: outcomes from our experience. *Breast*. 2008;17(6):596–603
 26. Neinstein LS, ed. *Adolescent Health Care*. 5th ed. Philadelphia, PA: Lippincott, Williams & Wilkins; 2008
 27. Rosen DS. Question from the clinician: adolescent gynecomastia. *Pediatr Rev*. 2003;24(9):317–319
 28. Lynne SD, Graber JA, Nichols TR, Brooks-Gunn J, Botvin GJ. Links between pubertal timing, peer influences, and externalizing behaviors among urban students followed through middle school. *J Adolesc Health*. 2007;40(2):181.e7–181.e13
 29. Graber JA, Seeley JR, Brooks-Gunn J, Lewinsohn PM. Is pubertal timing associated with psychopathology in young adulthood. *J Am Acad Child Adolesc Psychiatry*. 2004;43(6):718–726
 30. Rosen D, Foster C. Delayed puberty. *Pediatr Rev*. 2001;22(9):309–315
 31. Dreno B, Poli F. Epidemiology of acne. *Dermatology*. 2003;206(1):7–10
 32. Tan JK. Psychosocial impact of acne vulgaris: evaluating the evidence. *Skin Therapy Lett*. 2004;9(7):1–3, 9
 33. Jaccard J, Dodge T, Dittus P. Parent-adolescent communication about sex and birth control: a conceptual framework. *New Dir Child Adolesc Dev*. 2002;(97):9–41
 34. Dilorio C, Pluhar E, Belcher L. Parent-child communication about sexuality: a review of the literature from 1980–2002. *J HIV/AIDS Prev Educ Adolesc Child*. 2003;5(3/4):7–32
 35. Kirkman M, Rosenthal DA, Feldman SS. Talking to a tiger: fathers reveal their difficulties in communicating about sexuality with adolescents. *New Dir Child Adolesc Dev*. 2002;(97):57–74
 36. Martino SC, Elliott MN, Corona R, Kanouse DE, Schuster MA. Beyond the “big talk”: the roles of breadth and repetition in parent-adolescent communication about sexual topics. *Pediatrics*. 2008;121(3). Available at: www.pediatrics.org/cgi/content/full/121/3/e612
 37. American Academy of Pediatrics, Committee on Psychosocial Aspects of Child and Family Health and Committee on Adolescence. Sexuality education for children and adolescents. *Pediatrics*. 2001;108(2):498–502
 38. Henry J. Kaiser Family Foundation; Seventeen Magazine. *Sexual Health Care and Counsel: SexSmarts—A Series of National Surveys of Teens About Sex*. Menlo Park, CA: Henry J. Kaiser Family Foundation and Seventeen Magazine; 2001. Available at: www.kff.org/entpartnerships/upload/SexSmarts-Survey-Sexual-Health-Care-and-Counsel-Summary.pdf. Accessed June 22, 2010
 39. Ackard DM, Neumark-Sztainer D. Health care information sources for adolescents: age and gender differences on use, concerns, and needs. *J Adolesc Health*. 2001;29(3):170–176
 40. Ku LC, Sonenstein FL, Pleck JH. The association of AIDS education and sex education with sexual behavior and condom use among teenage men. *Fam Plann Perspect*. 1992;24(3):100–106
 41. Frappier JY, Kaufman M, Baltzer F, et al. Sex and sexual health: a survey of Canadian youth and mothers. *Paediatr Child Health*. 2008;13(1):25–30
 42. Brown JD, Strasburger VC. From Calvin Klein to Paris Hilton and MySpace: adolescents, sex, and the media. *Adolesc Med State Art Rev*. 2007;18(3):484–507
 43. Bell DL. Adolescent male sexuality. *Adolesc Med*. 2003;14(3):583–594, vi
 44. Beals KP, Peplau LA, Gable SL. Stigma management and well-being: the role of perceived social support, emotional processing, and suppression. *Pers Soc Psychol Bull*. 2009;35(7):867–879
 45. Elizur Y, Ziv M. Family support and acceptance, gay male identity formation, and psychological adjustment: a path model. *Fam Process*. 2001;40(2):125–144
 46. Frankowski BL; American Academy of Pediatrics, Committee on Adolescence. Sexual orientation and adolescents. *Pediatrics*. 2004;113(6):1827–1832
 47. Russell ST, Driscoll AK, Truong N. Adolescent same-sex romantic attractions and relationships: implications for substance use and abuse. *Am J Public Health*. 2002;92(2):198–202
 48. Russell ST, Franz BT, Driscoll AK. Same-sex romantic attraction and experiences of violence in adolescence. *Am J Public Health*. 2001;91(6):903–906
 49. Russell ST, Joyner K. Adolescent sexual orientation and suicide risk: evidence from a national study. *Am J Public Health*. 2001;91(8):1276–1281
 50. Russell ST, Seif H, Truong NL. School outcomes of sexual minority youth in the United States: evidence from a national study. *J Adolesc*. 2001;24(1):111–127
 51. Eccles TA, Sayegh MA, Fortenberry JD, Zimet GD. More normal than not: a qualitative assessment of the developmental experiences of gay male youth. *J Adolesc Health*. 2004;35(5):425.e411–425.e428
 52. Sorenson RC. *Adolescent Sexuality in Contemporary America*. New York, NY: World Publishing; 1973
 53. Leitenberg H, Detzer MJ, Srebnik D. Gender differences in masturbation and the relation of masturbation experience in preadolescence and/or early adolescence to sexual behavior and sexual adjustment in young adulthood. *Arch Sex Behav*. 1993;22(2):87–98
 54. Smith AM, Rosenthal DA, Reichler H. High schoolers masturbatory practices: their relationship to sexual intercourse and personal characteristics. *Psychol Rep*. 1996;79(2):499–509
 55. Janczewski Z, Bablok L. Semen characteristics in pubertal boys. I. Semen quality after first ejaculation. *Arch Androl*. 1985;15(2–3):199–205
 56. Martinez GM, Chandra A, Abma JC, Jones J, Mosher WD. Fertility, contraception, and fatherhood: data on men and women from cycle 6 (2002) of the 2002 National Survey of Family Growth. *Vital Health Stat*. 2006;(26):1–142
 57. Chandra A, Mosher WD, Copen C, Sioneau C. *Sexual Behavior, Sexual Attraction, and Sexual Identity in the United States: Data From the 2006–2008 National Survey of Family Growth*. Division of Vital Statistics, Centers for Disease Control and Prevention/National Center for Health Statistics and Division of HIV/AIDS Prevention, Centers for Disease Control and Prevention/National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention; 2011. Number 36
 58. Brewster KL, Tillman KH. Who’s doing it? Patterns and predictors of youths’ oral sexual experiences. *J Adolesc Health*. 2008;42(1):73–80
 59. Cornell JL, Halpern-Felsher BL. Adoles-

- cents tell us why teens have oral sex. *J Adolesc Health*. 2006;38(3):299–301
60. Halpern-Felsher BL, Cornell JL, Kropp RY, Tschann JM. Oral versus vaginal sex among adolescents: perceptions, attitudes, and behavior. *Pediatrics*. 2005; 115(4):845–851
 61. Mosher WD, Chandra A, Jones J. Sexual behavior and selected health measures: men and women 15–44 years of age, United States, 2002. *Adv Data*. 2005;(362):1–55
 62. Ericksen KP, Trocki KF. Behavioral risk factors for sexually transmitted diseases in American households. *Soc Sci Med*. 1992; 34(8):843–853
 63. Shafer MA, Hilton JF, Ekstrand M, et al. Relationship between drug use and sexual behaviors and the occurrence of sexually transmitted diseases among high-risk male youth. *Sex Transm Dis*. 1993;20(6): 307–313
 64. Ross MW, Essien EJ, Williams ML, Fernandez-Esquer ME. Concordance between sexual behavior and sexual identity in street outreach samples of four racial/ethnic groups. *Sex Transm Dis*. 2003;30(2): 110–113
 65. Pathela P, Hajat A, Schillinger J, Blank S, Sell R, Mostashari F. Discordance between sexual behavior and self-reported sexual identity: a population-based survey of New York City men [published correction appears in *Ann Intern Med*. 2006;145(12): 936]. *Ann Intern Med*. 2006;145(6): 416–425
 66. Henry J. Kaiser Family Foundation; Seventeen Magazine. *Gender Roles: SexSmarts—A Series of National Surveys of Teens About Sex*. Menlo Park, CA: Henry J. Kaiser Family Foundation and Seventeen Magazine; 2002. Available at: www.kff.org/entpartnerships/upload/Gender-Rolls-Summary.pdf. Accessed June 22, 2010
 67. National Campaign to Prevent Teen and Unplanned Pregnancy. Science says No. 6: the sexual attitudes and behavior of male teens. Available at: www.thenationalcampaign.org/resources/males.aspx. Accessed June 22, 2010
 68. National Campaign to Prevent Teen and Unplanned Pregnancy. With one voice: America's adults and teens sound off about teen pregnancy. Available at: www.thenationalcampaign.org/resources/pdf/pubs/WOV2007_fulltext.pdf. Accessed June 22, 2010
 69. Bensley LS, Van Eenwyk J, Simmons KW. Self-reported childhood sexual and physical abuse and adult HIV-risk behaviors and heavy drinking. *Am J Prev Med*. 2000;18(2): 151–158
 70. Metcalfe M, Oppenheimer R, Dignon A, Palmer RL. Childhood sexual experiences reported by male psychiatric patients. *Psychol Med*. 1990;20(4):925–929
 71. Rohsenow DJ, Corbett R, Devine D. Molested as children: a hidden contribution to substance abuse [published correction appears in *J Subst Abuse Treat*. 1988;5(2): 129]? *J Subst Abuse Treat*. 1988;5(1):13–18
 72. Holmes WC, Slap GB. Sexual abuse of boys: definition, prevalence, correlates, sequelae, and management. *JAMA*. 1998; 280(21):1855–1862
 73. Donovan P. Can statutory rape laws be effective in preventing adolescent pregnancy? *Fam Plann Perspect*. 1997;29(1): 30–34, 40
 74. Lewin Group. Statutory rape: a guide to state laws and reporting requirements. Available at: www.lewin.com/publications/Publication/75. Accessed October 5, 2011
 75. American Academy of Pediatrics, Council on Communications and Media. Policy statement: media education. *Pediatrics*. 2010;126(5):1012–1017
 76. Eaton DK, Kann L, Kinchen S, et al; Centers for Disease Control and Prevention. Youth risk behavior surveillance: United States, 2007. *MMWR Surveill Summ*. 2008;57(4): 1–131
 77. Ashley OS, Foshee VA. Adolescent help-seeking for dating violence: prevalence, sociodemographic correlates, and sources of help. *J Adolesc Health*. 2005; 36(1):25–31
 78. Zolotor AJ, Denham AC, Weil A. Intimate partner violence. *Prim Care*. 2009;36(1): 167–179
 79. Rhodes KV, Houry D, Cerulli C, Straus H, Kaslow NJ, McNutt LA. Intimate partner violence and comorbid mental health conditions among urban male patients. *Ann Fam Med*. 2009;7(1):47–55
 80. American Academy of Pediatrics, Committee on Injury, Violence, and Poison Prevention. Policy statement: role of the pediatrician in youth violence prevention. *Pediatrics*. 2009;124(1):393–402
 81. Alpert EJ, Sege RD, Bradshaw YS. Interpersonal violence and the education of physicians. *Acad Med*. 1997;72(1 suppl): S41–S50
 82. Greydanus DE, Pratt HD, Baxter T. Sexual dysfunction and the primary care physician. *Adolesc Med*. 1996;7(1):9–26
 83. Richters J, Grulich AE, de Visser RO, Smith AM, Rissel CE. Sex in Australia: sexual difficulties in a representative sample of adults. *Aust NZ J Public Health*. 2003;27(2): 164–170
 84. Weinstock H, Berman S, Cates W Jr. Sexually transmitted diseases among American youth: incidence and prevalence estimates, 2000. *Perspect Sex Reprod Health*. 2004;36(1):6–10
 85. Centers for Disease Control and Prevention. HIV surveillance report, 2009. 2011 vol 21. Available at: www.cdc.gov/hiv/topics/surveillance/resources/reports. Accessed October 5, 2011
 86. Institute of Medicine, Committee on Prevention and Control of Sexually Transmitted Diseases. *The Hidden Epidemic: Confronting Sexually Transmitted Diseases*. Eng TR, Butler WT, eds. Washington, DC: National Academies Press; 1997
 87. Miller WC, Ford CA, Morris M, et al. Prevalence of chlamydial and gonococcal infections among young adults in the United States. *JAMA*. 2004;291(18):2229–2236
 88. Datta SD, Sternberg M, Johnson RE, et al. Gonorrhea and chlamydia in the United States among persons 14 to 39 years of age, 1999 to 2002. *Ann Intern Med*. 2007; 147(2):89–96
 89. Miller WC, Swygard H, Hobbs MM, et al. The prevalence of trichomoniasis in young adults in the United States. *Sex Transm Dis*. 2005;32(10):593–598
 90. Cook LS, Koutsky LA, Holmes KK. Circumcision and sexually transmitted diseases. *Am J Public Health*. 1994;84(2):197–201
 91. Centers for Disease Control and Prevention. *Sexually Transmitted Disease Surveillance 2007*. Atlanta, GA: Centers for Disease Control and Prevention; 2008. Available at: www.cdc.gov/std/pubs [ordering information]. Accessed June 22, 2010
 92. Campsmith M, Rhodes P, Hall I. Estimated prevalence of undiagnosed HIV infection in the United States at the end of 2006 [abstr]. Presented at: 16th Conference on Retroviruses and Opportunistic Infections; February 8–11, 2009; Montreal, Quebec, Canada. Available at: www.retroconference.org/2009/Abstracts/33682.htm. Accessed June 22, 2010
 93. Valleroy LA, MacKellar DA, Karon JM, et al. HIV prevalence and associated risks in young men who have sex with men. Young Men's Survey Study Group. *JAMA*. 2000; 284(2):198–204
 94. US Preventive Services Task Force. Screening for HIV: recommendation statement. *Ann Intern Med*. 2005;143(1):32–37
 95. Weiss HA, Quigley MA, Hayes RJ. Male circumcision and risk of HIV infection in sub-

- Saharan Africa: a systematic review and meta-analysis. *AIDS*. 2000;14(15):2361–2370
96. Auvert B, Taljaard D, Lagarde E, Sobngwi-Tambekou J, Sitta R, Puren A. Randomized, controlled intervention trial of male circumcision for reduction of HIV infection risk: the ANRS 1265 Trial. *PLoS Med*. 2005;2(11):e298
 97. Bailey RC, Moses S, Parker CB, et al. Male circumcision for HIV prevention in young men in Kisumu, Kenya: a randomised controlled trial. *Lancet*. 2007;369(9562):643–656
 98. Gray RH, Kigozi G, Serwadda D, et al. Male circumcision for HIV prevention in men in Rakai, Uganda: a randomised trial. *Lancet*. 2007;369(9562):657–666
 99. Centers for Disease Control and Prevention. *CDC HIV/AIDS Science Facts: Male Circumcision and Risk for HIV Transmission and Other Health Conditions—Implications for the United States*. Atlanta, GA: Centers for Disease Control and Prevention; 2008. Available at: www.cdc.gov/hiv/resources/factsheets/PDF/circumcision.pdf. Accessed June 22, 2010
 100. Santelli JS, Orr M, Lindberg LD, Diaz DC. Changing behavioral risk for pregnancy among high school students in the United States, 1991–2007. *J Adolesc Health*. 2009;45(1):25–32
 101. Guttmacher Institute. *U.S. Teenage Pregnancy Statistics National and State Trends and Trends by Race and Ethnicity*. New York, NY: Guttmacher Institute; 2010. Available at: www.guttmacher.org/pubs/USTPtrends.pdf. Accessed January 1, 2011
 102. Finer LB, Henshaw SK. Disparities in rates of unintended pregnancy in the United States, 1994 and 2001. *Perspect Sex Reprod Health*. 2006;38(2):90–96
 103. Kirby D. *Emerging Answers 2007: Research Findings on Programs to Reduce Teen Pregnancy (Summary)*. Washington, DC: National Campaign to Prevent Pregnancy Prevention; 2001. Available at: www.thenationalcampaign.org/resources/pdf/pubs/EA2007_FINAL.pdf. Accessed June 22, 2010
 104. Gavin L, MacKay AP, Brown K, et al; Centers for Disease Control and Prevention. Sexual and reproductive health of persons aged 10–24 years: United States, 2002–2007. *MMWR Surveill Summ*. 2009;58(6):1–58
 105. Lindberg LD, Sonenstein FL, Ku L, Martinez G. Age differences between minors who give birth and their adult partners. *Fam Plann Perspect*. 1997;29(2):61–66
 106. Ventura SJ, Martin JA, Curtin SC, Mathews TJ. Report of final natality statistics, 1996. *Mon Vital Stat Rep*. 1998;46(11 suppl):1–99
 107. Landry DJ, Forrest JD. How old are U.S. fathers? *Fam Plann Perspect*. 1995;27(4):159–161, 165
 108. Darroch JE, Landry DJ, Oslak S. Age differences between sexual partners in the United States. *Fam Plann Perspect*. 1999;31(4):160–167
 109. Sonenstein F, Stewart K, Lindberg L, Pernas M, Williams S. *Involving Males in Preventing Teen Pregnancy: A Guide for Program Planners*. Washington, DC: Urban Institute; 1997
 110. Martin JA, Hamilton BE, Sutton PD, et al. Births: final data for 2005. *Natl Vital Stat Rep*. 2007;56(6):1–103
 111. Adelman WP, Joffe A. The adolescent with a painful scrotum. *Contemp Pediatr*. 2000;17(3):111
 112. Idahl A, Boman J, Kumlin U, Olofsson JI. Demonstration of *Chlamydia trachomatis* IgG antibodies in the male partner of the infertile couple is correlated with a reduced likelihood of achieving pregnancy. *Hum Reprod*. 2004;19(5):1121–1126
 113. Hillis SD, Joesoef R, Marchbanks PA, Wasserheit JN, Cates W Jr, Westrom L. Delayed care of pelvic inflammatory disease as a risk factor for impaired fertility. *Am J Obstet Gynecol*. 1993;168(5):1503–1509
 114. Hillis SD, Owens LM, Marchbanks PA, Amsterdam LF, Mac Kenzie WR. Recurrent chlamydial infections increase the risks of hospitalization for ectopic pregnancy and pelvic inflammatory disease. *Am J Obstet Gynecol*. 1997;176(1 pt 1):103–107
 115. Jones RB, Wasserheit JN. Introduction to the biology and natural history of sexually transmitted diseases. In: Wasserheit JN, Aral SO, Holmes KK, Hitchcock PJ, eds. *Research Issues in Human Behavior and Sexually Transmitted Disease in the AIDS Era*. Washington, DC: American Society for Microbiology; 1991:11–37
 116. Marchbanks PA, Annegers JF, Coulam CB, Strathy JH, Kurland LT. Risk factors for ectopic pregnancy: a population-based study. *JAMA*. 1988;259(12):1823–1827
 117. Goldenberg RL, Andrews WW, Yuan AC, MacKay HT, St Louis ME. Sexually transmitted diseases and adverse outcomes of pregnancy. *Clin Perinatol*. 1997;24(1):23–41
 118. Heath DT, McKenry PC, Leigh GK. The consequences of adolescent parenthood on men's depression, parental satisfaction, and fertility in adulthood. *J Soc Serv Res*. 1995;20(3–4):127–148
 119. Hoffman S. *By the Numbers: The Public Costs of Teen Childbearing*. Washington, DC: National Campaign to Prevent Teen Pregnancy; 2006. Available at: www.thenationalcampaign.org/resources/pdf/pubs/BTN_Full.pdf. Accessed June 22, 2010
 120. Fine J. Adolescent pregnancy prevention in managed care. *Womens Health Issues*. 1998;8(3):148–168
 121. Trussell J. Contraceptive efficacy. In: Hatcher RA, Trussell J, Nelson AL, Cates W, Stewart FH, Kowal D, eds. *Contraceptive Technology*. 19th Rev ed. New York, NY: Ardent Media; 2007:747–826
 122. American Academy of Pediatrics, Committee on Adolescence. Condom use by adolescents. *Pediatrics*. 2001;107(6):1463–1469
 123. Centers for Disease Control and Prevention. *Condoms and STDs: Fact Sheet for Public Health Personnel*. Atlanta, GA: Centers for Disease Control and Prevention; 2009. Available at: www.cdc.gov/condomeffectiveness/latex.htm. Accessed June 22, 2010
 124. Paz-Bailey G, Koumans EH, Sternberg M, et al. The effect of correct and consistent condom use on chlamydial and gonococcal infection among urban adolescents. *Arch Pediatr Adolesc Med*. 2005;159(6):536–542
 125. Crosby RA, Salazar LF, Yarber WL, et al. A theory-based approach to understanding condom errors and problems reported by men attending an STI clinic. *AIDS Behav*. 2008;12(3):412–418
 126. Crosby RA, Yarber WL, Sanders SA, et al. Men with broken condoms: who and why? *Sex Transm Infect*. 2007;83(1):71–75
 127. Crosby RA, Diclemente RJ, Wingood GM, et al. Correlates of condom failure among adolescent males: an exploratory study. *Prev Med*. 2005;41(5–6):873–876
 128. Crosby RA, DiClemente RJ, Wingood GM, et al. Condom failure among adolescents: implications for STD prevention. *J Adolesc Health*. 2005;36(6):534–536
 129. Manlove J, Ikramullah E, Terry-Humen E. Condom use and consistency among male adolescents in the United States. *J Adolesc Health*. 2008;43(4):325–333
 130. Dariotis JK, Sonenstein FL, Gates GJ, et al. Changes in sexual risk behavior as young men transition to adulthood. *Perspect Sex Reprod Health*. 2008;40(4):218–225
 131. Jones RK, Fennell J, Higgins JA, Blanchard K. Better than nothing or savvy risk-reduction practice? The importance of withdrawal. *Contraception*. 2009;79(6):407–410
 132. Merkh RD, Whittaker PG, Baker K, Hock-

- Long L, Armstrong K. Young unmarried men's understanding of female hormonal contraception. *Contraception*. 2009;79(3):228–235
133. Henry J. Kaiser Family Foundation; Seventeen Magazine. *Birth Control and Protection. SexSmarts—A Series of National Surveys of Teens About Sex*. Menlo Park, CA: Henry J. Kaiser Family Foundation and Seventeen Magazine; 2004. Available at: www.kff.org/entpartnerships/7106.cfm. Accessed June 22, 2010
 134. American Academy of Pediatrics, Committee on Adolescence. Emergency contraception. *Pediatrics*. 2005;116(4):1026–1035
 135. Manlove J, Ryan S, Franzetta K. Patterns of contraceptive use within teenagers' first sexual relationships. *Perspect Sex Reprod Health*. 2003;35(6):246–255
 136. Ryan S, Franzetta K, Manlove J, Holcombe E. Adolescents' discussions about contraception or STDs with partners before first sex. *Perspect Sex Reprod Health*. 2007;39(3):149–157
 137. Brückner H, Bearman P. After the promise: the STD consequences of adolescent virginity pledges. *J Adolesc Health*. 2005;36(4):271–278
 138. Frankel L. An appeal for additional research about the development of heterosexual male sexual identity. *J Psychol Human Sex*. 2004;16(4):1–16
 139. Marcell AV, Ford CA, Pleck JH, Sonenstein F. Masculine beliefs, parental communication, and adolescent males' health care use. *Pediatrics*. 2007;119(4). Available at: www.pediatrics.org/cgi/content/full/119/4/e966
 140. Goldstein LS, Chapin JL, Lara-Torre E, Schulkin J. The care of adolescents by obstetrician-gynecologists: a first look. *J Pediatr Adolesc Gynecol*. 2009;22(2):121–128
 141. Kalmuss D, Austrian K. Real men do... real men don't: young Latino and African American men's discourses regarding sexual health care utilization. *Am J Mens Health*. 2010;4(3):218–230
 142. Ford C, Best D, Miller W. Confidentiality and adolescents' willingness to consent to STD testing. *Arch Pediatr Adolesc Med*. 2001;155(9):1072–1073
 143. Fortenberry JD, McFarlane M, Bleakley A, et al. Relationships of stigma and shame to gonorrhea and HIV screening. *Am J Public Health*. 2002;92(3):378–381
 144. Lowery LM, Chung S, Ellen JM. Social support and sexually transmitted disease related healthcare utilisation in sexually experienced African-American adolescents. *Sex Transm Infect*. 2005;81(1):63–66
 145. Marcell AV, Howard TL, Plowden K, Watson C. Exploring women's perceptions about their role in supporting partners' and sons' reproductive health care. *Am J Mens Health*. 2010;4(4):297–304
 146. Elster AB, Marcell AV. Health care of adolescent males: overview, rationale, and recommendations. *Adolesc Med*. 2003;14(3):525–540, v
 147. Rosen DS, Neinstein LS. Preventive health care for adolescents. In: Neinstein LS, Gordon CM, Katzman DK, Rosen DS, Woods ER, eds. *Adolescent Health Care: A Practical Guide*. 5th ed. Philadelphia, PA: Lippincott, Williams & Wilkins; 2008:44–80
 148. Ford C, English A, Sigman G. Confidential health care for adolescents: position paper for the society for adolescent medicine. *J Adolesc Health*. 2004;35(2):160–167
 149. Guttmacher Institute. *State Policies in Brief: An Overview of Minors' Consent Law*. New York, NY: Guttmacher Institute; 2009. Available at: www.guttmacher.org/statecenter/spibs/spib_OMCL.pdf. Accessed June 22, 2010
 150. Branson BM, Handsfield HH, Lampe MA, et al. Centers for Disease Control and Prevention. Revised recommendations for HIV testing of adults, adolescents, and pregnant women in health-care settings. *MMWR Recomm Rep*. 2006;55(RR-14):1–17; quiz CE1–CE4
 151. Centers for Disease Control and Prevention. *Male Chlamydia Screening Consultation*. Atlanta, GA: Centers for Disease Control and Prevention; 2006. Available at: www.cdc.gov/std/chlamydia/ChlamydiaScreening-males.pdf. Accessed June 22, 2010
 152. Workowski KA, Berman S. Sexually transmitted diseases treatment guidelines, 2010 [published correction appears in *MMWR Recomm Rep*. 2011;60(1):18]. *MMWR Recomm Rep*. 2010;59(RR-12):1–110
 153. US Preventive Services Task Force. *Behavioral Counseling to Prevent Sexually Transmitted Infections*. Rockville, MD: Agency for Healthcare Research and Quality; 2008. Available at: www.ahrq.gov/clinic/uspstf/uspstfstds.htm. Accessed June 22, 2010
 154. US Preventive Services Task Force. Screening for HIV: recommendation statement. *Am Fam Physician*. 2005;72(11):2287–2292
 155. US Preventive Services Task Force. *Screening for Gonorrhea*. Rockville, MD: Agency for Healthcare Research and Quality; 2005. Available at: www.ahrq.gov/clinic/uspstf/uspstfgono.htm. Accessed June 22, 2010
 156. US Preventive Services Task Force. Screening for chlamydial infection: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med*. 2007;147(2):128–134
 157. Calonge N. Screening for syphilis infection: recommendation statement. *Ann Fam Med*. 2004;2(4):362–365
 158. American Academy of Pediatrics, Committee on Infectious Diseases. Policy statement: recommended childhood and adolescent immunization schedules—United States, 2011. *Pediatrics*. 2011;127(2):387–388
 159. Advisory Committee on Immunization Practices. ACIP recommends all 11-12 year-old males get vaccinated against HPV. Available at: www.cdc.gov/media/releases/2011/a1025_ACIP_HPV_Vote.html. Accessed October, 25, 2011
 160. Solberg LI, Nordin JD, Bryant TL, Kristensen AH, Maloney SK. Clinical preventive services for adolescents. *Am J Prev Med*. 2009;37(5):445–454
 161. Schuster MA, Bell RM, Petersen LP, Kanouse DE. Communication between adolescents and physicians about sexual behavior and risk prevention. *Arch Pediatr Adolesc Med*. 1996;150(9):906–913
 162. Marcell AV, Lindberg LD. Females do not fare much better than males: prevalence of heterosexually active young peoples sexual/reproductive health care receipt; NSFG 2006-8. *J Adolesc Health*. 2010;46(2 suppl):S23–S24
 163. Turner CF, Ku L, Rogers SM, Lindberg LD, Pleck JH, Sonenstein FL. Adolescent sexual behavior, drug use, and violence: increased reporting with computer survey technology. *Science*. 1998;280(5365):867–873
 164. Paperny DM. Office-based computerized risk-assessment and health-education systems. *Adolesc Med State Art Rev*. 2007;18(2):256–270
 165. Paperny DM, Hedberg VA. Computer-assisted health counselor visits: a low-cost model for comprehensive adolescent preventive services. *Arch Pediatr Adolesc Med*. 1999;153(1):63–67
 166. Knight JR, Shrier LA, Bravender TD, Farrell M, Vander Bilt J, Shaffer HJ. A new brief screen for adolescent substance abuse. *Arch Pediatr Adolesc Med*. 1999;153(6):591–596
 167. Goldenring JM, Cohen E. Getting into adolescents heads. *Contemp Pediatr*. 1988;5(7):75–90
 168. Marcell AV. Making the most of the adolescent male health visit. Part 2: the

- physical exam. *Contemp Pediatr*. 2006; 23(6):38–46
169. Adelman WP, Joffe A. The adolescent male genital examination: what's normal and what's not. *Contemp Pediatr*. 1999;16(7): 76–92
 170. US Preventive Services Task Force. Screening for testicular cancer: recommendation statement. Available at: www.ahrq.gov/clinic/3rduspstf/testicular/testiculup.htm. Accessed May 1, 2011
 171. Lin K, Sharangpani R. Screening for testicular cancer: an evidence review for the U.S. Preventive Services Task Force. *Ann Intern Med*. 2010;153(6):396–399
 172. American Cancer Society. Detailed guide: testicular cancer—can testicular cancer be found early? Available at: www.cancer.org/Cancer/TesticularCancer/DetailedGuide/testicular-cancer-detection. Accessed January 23, 2011
 173. American Academy of Pediatrics, Committee on Practice and Ambulatory Medicine. The use of chaperones during the physical examination of the pediatric patient. *Pediatrics*. 1996;98(6 pt 1):1202
 174. Foxman B. Epidemiology of urinary tract infections: incidence, morbidity, and economic costs. *Am J Med*. 2002;113(suppl 1A):5S–13S
 175. Goldie SJ, Kuntz KM, Weinstein MC, Freedberg KA, Palefsky JM. Cost-effectiveness of screening for anal squamous intraepithelial lesions and anal cancer in human immunodeficiency virus-negative homosexual and bisexual men. *Am J Med*. 2000; 108(8):634–641
 176. Goldie SJ, Kuntz KM, Weinstein MC, Freedberg KA, Welton ML, Palefsky JM. The clinical effectiveness and cost-effectiveness of screening for anal squamous intraepithelial lesions in homosexual and bisexual HIV-positive men. *JAMA*. 1999;281(19): 1822–1829
 177. Ferraris A, Narasimhan K, Rahman I, Jorge I. Anal pap smear in high-risk patients: a poor screening tool. *South Med J*. 2008; 101(11):1185–1186
 178. American Academy of Pediatrics. *Red Book: 2009 Report of the Committee on Infectious Diseases*. 28th ed. Elk Grove, IL: American Academy of Pediatrics; 2009. Available at: <http://aapredbook.aappublications.org>. Accessed October 5, 2011
 179. Centers for Disease Control and Prevention. Recommendations for partner services programs for HIV infection, syphilis, gonorrhea, and chlamydial infection. *MMWR Recomm Rep*. 2008;57(RR-9):1–83; quiz CE1–CE4
 180. Kamb ML, Fishbein M, Douglas JM Jr, et al. Efficacy of risk-reduction counseling to prevent human immunodeficiency virus and sexually transmitted diseases: a randomized controlled trial. Project RESPECT Study Group. *JAMA*. 1998; 280(13):1161–1167
 181. Ryan C. *Supportive Families, Healthy Children: Helping Families With Lesbian, Gay, Bisexual and Transgender Children*. San Francisco, CA: Marian Wright Education Institute, San Francisco State University; 2009. Available at: familyproject.sfsu.edu. Accessed June 1, 2010

Male Adolescent Sexual and Reproductive Health Care

Arik V. Marcell, Charles Wibbelsman, Warren M. Seigel and the Committee on Adolescence

Pediatrics; originally published online November 28, 2011;

DOI: 10.1542/peds.2011-2384

Updated Information & Services

including high resolution figures, can be found at:
<http://pediatrics.aappublications.org/content/early/2011/11/22/peds.2011-2384>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
<http://pediatrics.aappublications.org/site/misc/Permissions.xhtml>

Reprints

Information about ordering reprints can be found online:
<http://pediatrics.aappublications.org/site/misc/reprints.xhtml>

PEDIATRICS is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. PEDIATRICS is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2011 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 0031-4005. Online ISSN: 1098-4275.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

