

Continuing APRETUDE After Planned Missed Injection

Adherence to the injection dosing schedule is strongly recommended. Individuals who miss their Target Injection Date should be clinically reassessed, including HIV-1 testing, to ensure resumption of APRETUDE remains appropriate.

If your patient plans to miss their Target Injection Date by more than 7 days, daily oral cabotegravir can be prescribed for a duration of up to 2 months to replace 1 missed scheduled every-2-month injection of APRETUDE.†

The first dose of oral PrEP should be taken approximately 2 months after the last injection dose of APRETUDE.

Restart injections with APRETUDE on the day oral dosing completes or within 3 days.

How much time has passed since their missed Target Injection Date?



**≤1 month since missed
Target Injection Date**

- **Resume injections on final day of oral cabotegravir or within 3 days**
- Continue with every-2-month dosing schedule thereafter

**>1 month since missed
Target Injection Date**

- **Repeat initiation injections (2 injections 1 month apart) on the final day of oral cabotegravir or within 3 days**
- Continue with every-2-month dosing schedule thereafter

Continuing APRETUDE After Unplanned Missed Injection

Adherence to scheduled injection visits is important. Your patient missed their Target Injection Date by >7 days and did not plan for it by taking oral therapy.

Clinically reassess the patient to determine whether APRETUDE remains appropriate, and if so, confirm HIV-1–negative status prior to injection.

How much time has passed since their Target Injection Date?



**≤1 month since missed
Target Injection Date**

- **Resume injections as soon as possible**
- Continue with every-2-month dosing schedule thereafter

**>1 month since missed
Target Injection Date**

- **Restart initiation injections 1 month apart, for 2 consecutive months**
- Continue with every-2-month dosing schedule thereafter